



Screening (CMS Policy)	CPT/HCPCS	ICD-10 Codes	Medicare Covers	Frequency
Cardiovascular Disease Screening (NCD 190.23)	80061 – Lipid Panel (Cholesterol 82465, HDL 83718, Triglycerides 84478)	Z13.6 ¹	Part B patients without cardiovascular disease symptoms	
Diabetes Screening	82947, 82950, 82951, 83036 – Glucose, Tolerance, HbA1C Tests	Z13.1 ¹	Part B patients with diabetes risk factors (not previously diagnosed)	
Prostate Cancer Screening (NCD 210.1)	G0102 – Digital Rectal Exam; G0103 – PSA Test	Z12.5 ¹	Men aged 50+ with Part B. Coverage starts the day after 50th birthday	
Cervical Cancer Screening with HPV Test (NCD 210.2.1)	G0476 – HPV Test (high-risk types)	Z11.51 ¹ and either Z01.411 or Z01.419	Asymptomatic women aged 30–65 with Part B	
Colorectal Cancer Screening (NCD 210.3)	G0104, G0105, G0121, G0327, G0328, 00812, 74263, 81528, 82270, 0464U, 0537U	See the current list of ICD-10 codes in the 210.3 Colorectal Cancer Screening coding file ¹	Part B patients aged 45–85, asymptomatic and average risk	
Hepatitis B Screening (NCD 210.6)	G0499, 86704, 86706, 87340, 87341 – HBV Panel and Antigen Tests	*People with ESRD: Z11.59 and N18.6¹ *Asymptomatic, Non-Pregnant Adolescents and High-Risk Adults: Z11.59 and Z72.89^{1,2} *Asymptomatic, Non-Pregnant Adolescents and Adults, Subsequent Visits: Z11.59 and one of the following, F11.10 – F11.988, F13.10 – F13.988, F14.10 – F14.988, F15.10 – F15.988, Z20.2, Z20.5, Z72.52 or Z72.53¹ *Pregnant Women: Z11.59 and 1 of these: Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93^{1,3} *High Risk Pregnant Women: Z11.59 and Z72.89 and 1 of these: O09.90, O09.91, O09.92, O09.931	Patients with Medicare Part B who are either: *High-risk, asymptomatic non-pregnant adolescents and adults *Pregnant Women	*Annually for patients with continued high risk who don't get HBV *One asymptomatic screening for non-pregnant adolescents and adults who meet the high-risk definition *One screening for pregnant women at first prenatal visit, and rescreening at delivery for those with new or continued risk factors Note: This includes screening during the first prenatal visit for future pregnancies, even if the patient previously got the HBV shot or had a negative hepatitis B surface antigen screening result

See footnotes on page 3.

Page 1 of 3

Disclaimer: This associated condition reference guide is provided as an aid to physicians and office staff in determining when testing is medically necessary. The diagnosis must be applicable to the patient's symptoms or conditions and must be consistent with documentation in the patient's medical record. Diagnostic Laboratory of Oklahoma (DLO) does not recommend any diagnosis codes or conditions and will only submit diagnosis information provided to us by the ordering physician or his/her designated staff. The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the billed payor.

DLOLab.com DLO®, Diagnostic Laboratory of Oklahoma®, any associated logos, and all associated Diagnostic Laboratory of Oklahoma/DLO registered or unregistered trademarks are the property of DLO. All third-party marks—® and —are the property of their respective owners. © 2026 Diagnostic Laboratory of Oklahoma, LLC. All rights reserved.

Screening (CMS Policy)	CPT/HCPCS	ICD-10 Codes	Medicare Covers	Frequency
Hepatitis C Screening (NCD 210.13)	G0472, G0567 – HCV Antibody or DNA or RNA Test	Z11.59, Z72.89, F19.20 ^{1, 2}	Adults with Part B, high risk or born 1945–1965. Had blood transfusion before 1992	Once for patients born 1945–1965. An initial screening for high-risk patients who had a blood transfusion before 1992 and patients with current or history of illicit injection drug use
HIV Screening (NCD 210.7)	G0432, G0433, G0435, G0475, 80081 – HIVObstetric Panels	Increased risk factors not reported Z11.4, Increased risk factors reported Z11.4 & ² (see below). Pregnant patients ³ (see below) ¹	Part B patients without regard to perceived risk or at increased HIV risk, including anyone who asks for the test or pregnant women	Annually for patients 15–65 without regard to risk. Annually for patients younger than 15 and adults over 65 at increased risk. Pregnant patients, 3 times per pregnancy (1 when diagnosed as pregnant, 2 during 3rd trimester, 3 at labor, if their clinician orders it)
Screening Pap Test (NCD 210.2)	G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091 – Pap Smear Tests	High risk: Z72.51 – Z72.53, Z77.29, Z77.9, Z91.89, Z92.850, Z92.858, Z92.86, Z92.89 Low risk: Z01.411, Z01.419, Z12.4, Z12.72, Z12.79, Z12.89 ¹	Female patients with Medicare part B	Annually (or 11 months after the month of the last covered exam) for women at high risk for developing cervical or vaginal cancer or childbearing age with abnormal Pap test within past 3 years. Or every 2 years (or 23 months after the month of the last covered exam for low-risk women)

Screening (CMS Policy)	CPT/HCPCS	ICD-10 Codes	Medicare Covers	Frequency
STI Screening & HIBC (NCD 210.10)	Chlamydia - 86631, 86632, 87110, 87270, 87320, 87490, 87491, 87810, 87800 Gonorrhea - 87590, 87591, 87850, 87800 Chlamydia and Gonorrhea - 0402U and 0455U Syphilis - 86592, 86593, 86780 Hepatitis B Surface Antigen - See the Hepatitis B screening information above	Non-Pregnant and High-Risk Women (screening for chlamydia, gonorrhea, or syphilis: Z11.3 and one of these Z72.51, Z72.52, Z72.53, Z72.89 ^{1,3} High-Risk Pregnant Women (screening for chlamydia, gonorrhea or syphilis) Z11.3 and one of these: Z72.51, Z72.52, Z72.53, Z72.89 and 1 of these: Z34.00 - Z34.03, Z34.80-Z34.83, Z34.90 - Z34.93, O09.90-O09.931 Pregnant Women (Screening for Syphilis) Z11.3 and 1 on these Z34.00 - Z34.03, Z34.80 - Z34.83, Z34.90 - Z34.93, O09.90 - O09.931 High-Risk Men (screening for Syphilis) Z11.3 and any of these Z72.51, Z72.52, Z72.53, Z72.891	Part B patients who are sexually active adolescents/ adults at increased STI risk or primary care provider refers them to a Medicare-eligible primary care provider in a primary care setting	*Chlamydia, gonorrhea, and syphilis in women at increased risk but not pregnant, 1 screening per year *Syphilis in men at increased risk, 1 per year *Up to two, 20 – 30-minute, annual face-to-face HIBC sessions for all sexually active adolescents, and for adults at increased risk Up to 2 screenings for chlamydia and gonorrhea per pregnancy for women at increased STI risk and continued increased risk for second screening One syphilis screening per pregnancy for pregnant women (up to 2 additional screenings, 1 in 3rd trimester and 1 at delivery if the patient is at continued increased risk for STIs)

Footnotes

¹ Additional ICD-10 codes may apply. Find individual change requests and specific ICD-10-CM service codes we cover on [CMS ICD-10](#). Find your [MAC's website](#) for information.

² High Risk ICD-10 Codes: Z72.51, Z72.52, Z72.53 & Z72.89

³ OB Codes: Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, O09.90, O09.91, O09.92, or O09.93

To view the complete policy, please refer to the BCBSTX website referenced below:

<https://www.cms.gov/Medicare/Prevention/PrevntionGenInfo/medicare-preventive-services/MPS-QuickReferenceChart-1.html>

Last Updated: 7/16/2026